## Math 223a: Algebraic Number Theory Fall 2019

Homework #1

due Thursday, September 12 at noon

**Problem 1.** Show that every subgroup of  $\hat{\mathbb{Z}}$  of finite index is open.

**Problem 2.** We call an algebraic field extension L|K abelian if it is a Galois extension with abelian Galois group.

Let M|K be a Galois extension with Galois group G. Show that M|K has a (unique) maximal abelian subextension T|K: any subextension L|K of M|K is abelian if and only if  $L \subseteq T$ .

Show that  $\operatorname{Gal}(M|T) = \overline{[G,G]}$  is the topological closure of the commutator subgroup of G.

**Problem 3.** Let  $K = \mathbb{Q}(\sqrt{2}, \sqrt{3}, \sqrt{5}, \dots)$  be the smallest field extension of  $\mathbb{Q}$  containing the square roots of all prime numbers.

- a) Show that  $\operatorname{Gal}(K|\mathbb{Q}) \cong \prod_{k=1}^{\infty} \mathbb{Z}/2\mathbb{Z}$ , with the product topology (obtained from the discrete topology on  $\mathbb{Z}/2\mathbb{Z}$ ). How does the element of  $\operatorname{Gal}(K|\mathbb{Q})$  corresponding to a tuple  $(a_k)_k \in \prod_{k=1}^{\infty} \mathbb{Z}/2\mathbb{Z}$  act on K?
- b) Show that  $\operatorname{Gal}(K|\mathbb{Q})$  has a subgroup H of finite index which is not open.

**Problem 4.** Recall some notation: We say that two sets A and B have the same cardinality (written as |A| = |B|) if there is a bijection  $A \xrightarrow{\sim} B$ . We say that the cardinality of A is at most the cardinality of B (written as  $|A| \leq |B|$ ) if there is an injection  $A \hookrightarrow B$ . This is equivalent to the existence of a surjection  $B \to A$ . We also know that  $|A| \leq |B|$  and  $|B| \leq |A|$  implies that |A| = |B|. For example,  $|\mathbb{N}| < |\mathbb{R}| = |2^{\mathbb{N}}|$  where  $2^{\mathbb{N}}$  denotes the set of subsets of  $\mathbb{N}$ . A set A is countable if and only if  $|A| \leq |\mathbb{N}|$ . (We know that  $\overline{\mathbb{Q}}$  is countable because  $\overline{\mathbb{Q}} = \bigcup_{f(X) \in \mathbb{Q}[X]} \{\alpha \in \overline{\mathbb{Q}} \mid f(\alpha) = 0\}$  is the union of countably many countable (in fact finite) sets.)

- a) Show that  $|\operatorname{Gal}(\overline{\mathbb{Q}}|\mathbb{Q})| = |2^{\mathbb{N}}|.$
- b) Show that  $|\{H \subseteq \text{Gal}(\overline{\mathbb{Q}}|\mathbb{Q}) \text{ open subgroup}\}| = |\mathbb{N}|.$

- c) Show that  $|\{H \subseteq \operatorname{Gal}(\overline{\mathbb{Q}}|\mathbb{Q}) \text{ closed subgroup}\}| = |2^{\mathbb{N}}|.$
- d) (bonus) What is  $|\{H \subseteq \operatorname{Gal}(\overline{\mathbb{Q}}|\mathbb{Q}) \text{ subgroup}\}|$ ?